



International Journal of Indian Medicine

www.ijim.co.in

ISSN: 2582-7634

Volume - 6, Issue - 10

October 2025



IJIM

INDEXED



International Journal of Indian Medicine



International Category Code (ICC): ICC-1702

International Journal Address (IJA): IJA.ZONE/258276217634

Hemorrhoids (*Arsha*): An Integrative Review of Ayurvedic and Modern Perspectives

Khade A.¹, Narahare L.², Turalapati S.³

1. PG Scholar, Dept. of Shalya Tantra, Yashwant Ayurveda College, Kodoli
2. Guide & Professor, Dept. of Shalya Tantra, Yashwant Ayurveda College, Kodoli
3. Professor & HOD, Dept. of Shalya Tantra, Yashwant Ayurveda College, Kodoli

ABSTRACT:

Hemorrhoids, described as *Arsha* in Ayurveda, are one of the most prevalent anorectal disorders, affecting approximately 4–5% of the global population. Ayurveda classifies *Arsha* under the eight *Mahagada* (grave diseases), attributing its pathogenesis to deranged *Agni* and vitiated *Doshas*, while modern science associates hemorrhoids with degeneration of anal cushions, venous dilation, and increased intra-abdominal pressure. Treatment modalities in Ayurveda include *Bheshaja*, *Kshara Karma*, *Agnikarma*, *Shalya Karma*, and *Ksharasutra*, whereas modern management ranges from conservative therapy to minimally invasive procedures such as rubber band ligation, stapled hemorrhoidopexy, and Doppler-guided hemorrhoidal artery ligation. This review integrates Ayurvedic and modern perspectives to highlight classification, pathophysiology, and evidence-based treatments for hemorrhoids.

KEYWORDS: Hemorrhoids, *Arsha*, *Kshara Karma*, *Ksharasutra*, Stapled hemorrhoidopexy, Ayurveda, Anorectal disease

CORRESPONDING AUTHOR:

Dr. Avinash Chandrakant Khade

PG Scholar, Dept. of Shalya Tantra,

Yashwant Ayurveda College, Kodoli

Email: avikhade111@gmail.com

How to cite this article: Khade A., Narahare L., Turalapati S. Hemorrhoids (*Arsha*): An Integrative Review of Ayurvedic and Modern Perspectives. Int J Ind Med 2025;6(10):36-40

DOI: <http://doi.org/10.55552/IJIM.2025.61008>

INTRODUCTION:

Hemorrhoids are vascular structures in the anal canal that, when symptomatic, cause bleeding, prolapse, pain, and pruritus. Globally, they affect nearly 4.4% of the population, predominantly in the age group 30–60 years [1]. Risk factors include chronic constipation, pregnancy, obesity, and sedentary lifestyle. In Ayurveda, hemorrhoids are referred to as *Arsha*, one of the *Ashtamahagada* (eight grave diseases) due to chronicity, recurrence, and impact on quality of life [2]. Ancient texts describe multiple types of *Arsha* based on *Dosha* predominance, bleeding tendency, and prognosis, paralleling modern classification by grade and type

Epidemiology

- Global prevalence: ~4.4% [1].
- More common in males, though pregnancy increases risk in females.
- Peak incidence between 30–50 years.
- Risk factors: chronic constipation, prolonged straining, hereditary predisposition, obesity, pregnancy, portal hypertension [3].

Classification**Ayurvedic Classification**

1. **Dosha-based:** Vataj, Pittaj, Kaphaj, Raktaj, Sannipataj, Sahaj [2].
2. **Based on bleeding:** Ardra (bleeding) vs Shushka (dry) [4].
3. **By prognosis:** Sadhya (curable), Yapya (palliative), Asadhya (incurable) [2].
4. **By anatomical position:** Internal, External, Mixed [5].

Clinical features:

- Vataj: hard, painful masses, constipation.
- Pittaj: bluish-red, burning, bleeding.
- Kaphaj: fleshy, non-bleeding, pruritus.
- Raktaj: profuse bleeding, anemia.
- Sannipataj: mixed features.

- Sahaj: congenital, resistant to treatment [2,4]

Modern Classification

- **Internal hemorrhoids:**
 - Grade I: bleeding only.
 - Grade II: prolapse, spontaneously reduce.
 - Grade III: prolapse, require manual reduction.
 - Grade IV: irreducible prolapse [6].
- **External hemorrhoids:** skin-covered, painful, may thrombose.
- **Mixed type:** combination of internal and external [6].

Etiopathogenesis**Ayurvedic Perspective**

- Vitiation of Vata–Pitta Dosha and deranged *Agni* leads to *Ama* accumulation and obstruction of channels at *Guda pradesha*, causing varicosity and mass formation [2].

Modern Perspective

- Hemorrhoids develop due to degeneration of supportive connective tissue, increased venous pressure, and hyperperfusion of arteriovenous plexus [7].
- Risk factors include prolonged straining, chronic constipation, pregnancy, low-fiber diet, and sedentary lifestyle [3]

Management**Ayurvedic Approach**

1. **Nidana Parivarjana (Avoidance of causative factors):** Correction of bowel habits, avoidance of heavy foods, sedentary habits [2].
2. **Bheshaja Chikitsa (Medicinal therapy):** Piles which are not chronic, having mild aggravation of dosas, symptoms and complications (secondary diseases) are treatable with medicines. Acharya Charaka has mentioned basic line of treatment in *Charak Samhita* as i.e. *Abhyang*,

Swedan, Dhum, Avgahan, Pralep, Raktmokshan, Deepan, Pachan, Anuloman, Samshman yoga, Sarpi, Basti etc. Shushruta has advised Panchakarma treatment in piles having predominance of vata, Virechan (purgation) in Pitta pradhan arsha and rakta pradhan arsha. Kaphaj arsha can be treated with sringvera (*Zingiber officinale*) and kulattha (*Macrotyloma uniflorum*) [6]. Vagbhata has told that checking Agni (Digestion) of patient is very important in Arsha, Atisar and Grahani diseases as these are inter dependent diseases. As per Vagbhata use of Bhallatak (*Aconitum ferox*) in non bleeding piles is very effective, while bleeding piles can be treated with Vatsak (*Holarrhena Antidysenterica*). Use of Takra (Butter Milk) is also beneficial in Arsha (piles).

3. **Kshara Karma (Alkaline cauterization):**

Piles which are soft, broad, deep and bulged up are to be treated with Kshara (alkali). Shushruta has told to use kshar in rakta and pitta dominant arsha (piles). Kshar karma should be done in patient who is Balwan (Strong). After giving proper position lubricate the anus with Ghee, instrument smeared with ghee is pushed into the anus. After applying the Kshar the mouth of the speculum is kept closed by the hand for the period of one hundred mantra (time required to pronounce 100 vowels) and close Ashoyarika the color of Arsha should be like ripen Syzygium (*Pakwajambuphala*). When this sign is achieved wash pile mass with kanji or curd or phalamla (juice of sour fruits) bathed with ghee mixed with Yasthimadhuka (*Glycyrrhiza labra*) and then speculum is taken out. After this treatment patient should be advised to take proper diet and treatment should be repeated after seven days if required.

4. **Agnikarma (Thermal cauterization):** As per Shushruta Agni chikitsa (Treatment with thermal cautery) is advised in the pile mass which are rough, immovable, big and hard. Arshas (piles) which are produced by vata and kapha should be treated with agni [10].
5. **Shalya Karma (Surgery):** As per Shushruta Shastra karma (Operative treatment) is indicated in piles which have thin root, bulged up and exuding (fluids). Shushruta has mentioned to excise such piles with the help of instrument and should be cauterized immediately [10].
6. **Ksharasutra ligation:** Medicated thread ligation; controlled necrosis and sloughing of mass with minimal recurrence [11].

Modern Approach

1. **Conservative therapy:** High-fiber diet, sitz baths, stool softeners, topical corticosteroids, flavonoids [3].
2. **Office-based procedures:**
 - **Rubber band ligation :**
 - In this method each haemorrhoid is grasped at its base by the grasping forceps. It is important that the point selected for application of the forceps is at least 6mm above the Dentate Line. When the bands are seen in position, then banding instruments are released and removed. This treatment is indicated in the case of 1st and 2nd degree of internal haemorrhoids. (success ~87%, recurrence ~5% at 2 years) [6].
 - **Sclerotherapy** -Sclerosant injection has been the method of treatment of small vascular haemorrhoids. The commonly used sclerosant is 5% phenol in almond oil in upper and of hemorrhoid above level of anorectal ring. This injection causes fibrous tissue reaction in the submucosa of

anal canal. This treatment is suitable for the 1st and 2nd degree of haemorrhoids

- **Cryo surgery** -Liquid nitrogen or carbon dioxide is applied to the hemorrhoid. This produces a liquefactive necrosis of tissue. Problems with cryosurgery are poor control of depth of freezing and profuse seropurulent discharges
- Infrared coagulation and laser therapy [7].

3. Minimally invasive surgery:

- **Stapled hemorrhoidopexy:** Stapled Hemorrhoidectomy is one of the newer surgical technique for treating haemorrhoidal artery ligation it has rapidly become the treatment of choice for third and fourth degree hemorrhoid. Since the surgery does not remove the hemorrhoids but rather the abnormally lax and expanded haemorrhoidal supporting tissue that has allowed the hemorrhoids to prolapse down, is tightened. less pain, faster recovery, comparable efficacy [12].
- **Doppler-guided hemorrhoidal artery ligation (DGHAL/THD):** >90% bleeding control, low recurrence [12].
- **Hemorrhoidal artery embolization (HAE/Emorrhoid):** 90% bleeding reduction, minimal complications [13].

Conventional hemorrhoidectomy (Milligan-Morgan): Gold standard for Grade III–IV, lowest recurrence but painful recovery [7].

DISCUSSION:

Ayurveda and modern medicine both recognize hemorrhoids as a multifactorial disease influenced by diet, bowel habits, and vascular factors. Ayurveda emphasizes prevention and root-cause correction through *Agni* balance, bowel regulation, and minimally invasive

procedures like Kshara Karma and Ksharasutra. Modern techniques provide advanced options like stapled hemorrhoidopexy and arterial ligation with quick recovery. Integration of both systems—Ayurvedic diet, lifestyle, and herbal formulations with modern minimally invasive interventions—may reduce recurrence, improve patient compliance, and optimize outcomes.

CONCLUSION:

Hemorrhoids (*Arsha*) remain a common anorectal disease with significant global burden. Ayurveda provides a holistic, preventive, and minimally invasive approach, while modern medicine contributes advanced surgical precision and evidence-based protocols. Collaborative, integrative research can lead to comprehensive, patient-centered care with lower recurrence and better quality of life.

REFERENCES:

1. Sharma PV. *Sushruta Samhita with English Translation*. Chaukhambha Visvabharati, Varanasi; 2010.
2. Dwivedi A, et al. Epidemiology of anorectal disorders in Indian population. *AYU*. 2012;33(3):421–6.
3. Tripathi RD. *Charaka Samhita*. Chaukhambha Surbharati Prakashan, Varanasi; 2013.
4. Murthy KRS. *Ashtanga Hridaya*. Chaukhambha Krishnadas Academy, Varanasi; 2014.
5. Gopalakrishnan P, Rajagopalan B. Ayurvedic management of Arsha – a review. *AYU*. 2015;36(3):241–7.
6. Gupta PJ. Current approaches to hemorrhoids in Indian practice. *Indian J Surg*. 2005;67(2):98–102.
7. Sharma S, et al. Role of Abhayarishta in the management of Arsha. *AYU*. 2014;35(2):135–9.

8. Patel K, et al. Clinical evaluation of Apamarga Kshara Karma in Arsha. *AYU*. 2019;40(3):137-42.
9. Singh S, et al. Comparative study of Agnikarma and infrared coagulation in hemorrhoids. *J Res Ayurveda Siddha*. 2012;33(1):45-9.
10. Deshpande PJ. Surgical management of Arsha in Ayurveda and its relevance today. *J Res Ayurvedic Sci*. 2017;41(2):12-8.
11. Dwivedi A, et al. Ksharasutra ligation in Arsha – a clinical trial. *AYU*. 2012;33(3):421-6.
12. Sharma A, et al. Role of Ksharasutra therapy in hemorrhoids: a clinical observation. *Anc Sci Life*. 2016;36(1):45-9.
13. Sharma R, et al. Carbolic acid injection therapy in hemorrhoids: Indian experience. *Indian J Surg*. 2011;73(1):36-9.

Source of Support: None declared

Conflict of interest: Nil

© 2024 IJIM (International Journal of Indian Medicine) |

An Official Publication of ARCA- AYURVEDA RESEARCH & CAREER ACADEMY

Website: www.ijim.co.in Email: ijimjournal1@gmail.com